

The human monocytic cell line THP-1 was grown in a synthetic cell culture medium RPMI 1640 supplemented with 10 % foetal calf serum, 6mM glutamine and Penicillin-Streptomycin (at 50 µg streptomycin/ml, Gibco BRL). THP-1 cells were washed in HBSS (lacking Ca<sup>2+</sup> and Mg<sup>2+</sup>) + 1 mg/ml BSA and resuspended in the same buffer at a density of 3 x 10<sup>6</sup> cells/ml. The cells were then loaded with 1mM FURA-2/AM for 30 min at 37°C, washed twice in HBSS, and resuspended at 1x10<sup>6</sup> cells/ml. THP-1 cell suspension (0.9 ml) was added to a 5 ml disposable cuvette containing a magnetic stirrer bar and 2.1 ml of prewarmed (37°C) HBSS containing 1 mg/ml BSA, 1 mM MgCl<sub>2</sub> and 2 mM CaCl<sub>2</sub>. The cuvette was placed in a fluorescence spectrophotometer (Perkin Elmer, Norwalk, CT) and preincubated for 4 min at 37°C with stirring. Fluorescence was recorded over 70 sec and cells were stimulated by addition of hMCP-1 to the cuvette after 10 sec. [Ca<sup>2+</sup>]<sub>i</sub> was measured by excitation at 340 nm and 380 nm alternately and subsequent measurement of the intensity of the fluorescence emission at 510 nm. The ratio of the intensities of the emitted fluorescent light following excitation at 340 nm and 380 nm, (R), was calculated and displayed to give and estimate of cytoplasmic [Ca<sup>2+</sup>]<sub>i</sub> according to the equation:-

$$[Ca^{2+}]_i = K_d \frac{(R - R_{min})}{(R_{max} - R)} \left( \frac{Sf2/Sb2}{Sf2/Sb2} \right)$$

where the K<sub>d</sub> for FURA-2 Ca<sup>2+</sup> complex at 37°C was taken to be 224nm. R<sub>max</sub> is the maximal fluorescence ratio determined after addition of 10 mM Ionomycin, R<sub>min</sub> is the minimal ratio determined by the subsequent addition of a Ca<sup>2+</sup> free solution containing 5 mM EGTA, and Sf2/Sb2 is the ratio of fluorescence values at 380 nm excitation determined at R<sub>min</sub> and R<sub>max</sub>, respectively.

Stimulation of THP-1 cells with hMCP-1 induced a rapid, transient rise in [Ca<sup>2+</sup>]<sub>i</sub> in a specific and dose dependent manner. Dose response curves indicated an approximate EC<sub>50</sub> of 2 nm. Test compounds dissolved in DMSO (10µl) were assayed for inhibition of calcium release by adding them to the cell suspension 10 sec prior to ligand addition and measuring the reduction in the transient rise in [Ca<sup>2+</sup>]<sub>i</sub>. Test compounds were also checked for lack of agonist activity by addition in place of hMCP-1.

### c) hMCP-1 and RANTES mediated chemotaxis.

*In vitro* chemotaxis assays were performed using the human monocytic cell line THP-1. Cell migration through polycarbonate membranes was measured by enumerating those passing through either directly by Coulter counting or indirectly by use of a

colourimetric viability assay measuring the cleavage of a tetrazolium salt by the mitochondrial respiratory chain (Scudiero D.A. *et al.* 1988, *Cancer Res.*, **48**, 4827-4833).

Chemoattractants were introduced into a 96-well microtitre plate which forms the lower well of a chemotaxis chamber fitted with a PVP-free 5  $\mu$ m poresize polycarbonate adhesive framed filter membrane (NeuroProbe MB series, Cabin John, MD 20818, USA) according to the manufacturer's instructions. The chemoattractant was diluted as appropriate in synthetic cell culture medium, RPMI 1640 (Gibco) or supplemented with 2 mM glutamine and 0.5% BSA, or alternatively with HBSS with  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  without Phenol Red (Gibco) plus 0.1% BSA. Each dilution was degassed under vacuum for 30 min and was placed (400  $\mu$ l) in the lower wells of the chamber and THP-1 cells ( $5 \times 10^5$  in 100  $\mu$ l RPMI 1640 + 0.5% BSA) were incubated in each well of the upper chamber. For the inhibition of chemotaxis the chemoattractant was kept at a constant submaximal concentration determined previously (1 nM MCP-1) and added to the lower well together with the test compounds dissolved in DMSO (final DMSO concentration < 0.05% v/v) at varying concentrations. The chamber was incubated for 2 h at 37°C under 5 %  $\text{CO}_2$ . The medium was removed from the upper wells which were then washed out with 200  $\mu$ l physiological saline before opening the chamber, wiping dry the membrane surface and centrifuging the 96-well plate at 600 g for 5 min to harvest the cells. Supernatant (150  $\mu$ l) was aspirated and 10  $\mu$ l of cell proliferation reagent, WST-1, {4-[3-(4-iodophenyl)-2-(4-nitrophenyl)-2H-5-tetrazolio]-1,3-phenyl disulfonate} plus an electron coupling reagent (Boehringer Mannheim, Cat.no. 1644 807) was added back to the wells. The plate was incubated at 37°C for 3 h and the absorbance of the soluble formazan product was read on a microtitre plate reader at 450 nm. The data was input into a spreadsheet, corrected for any random migration in the absence of chemoattractant and the average absorbance values, standard error of the mean, and significance tests were calculated. hMCP-1 induced concentration dependent cell migration with a characteristic biphasic response, maximal 0.5-1.0 nm.

In an alternative form of the above assay, fluorescently tagged cells can be used in order to assist in end point detection. In this case, the THP-1 cells used are fluorescently tagged by incubation in the presence of 5mM Calcein AM (Glycine, N,N'-[[3',6'-bis(acetyloxy)-3-oxospiro[isobenzofuran-1(3H),9'-[9H]xanthene]-2',7'-diyl]bis(methylene)]bis[N-[2-[(acetyloxy)methoxy]-2-oxoethyl]]-bis[(acetyloxy)methyl] ester: Molecular Probes) for 45 minutes in the dark. Cells are harvested by centrifugation and resuspended in HBSS

(without Phenol Red) with  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$  and 0.1% BSA. 50 $\mu\text{l}$  ( $2 \times 10^5$  cells) of the cell suspension are placed on the filter above each well and, as above, the unit is incubated at 37°C for 2 hours under 5%  $\text{CO}_2$ . At the end of the incubation, cells are washed off the upper face of the filter with phosphate buffered saline, the filter removed from the plate and the number of cells attracted to either the underside of the filter or the lower well estimated by reading fluorescence at 485nm excitation, 538nm emission wavelengths (fmax, Molecular Devices). The data was input into a spreadsheet, corrected for any random migration in the absence of chemoattractant and the average fluorescence values, standard error of the mean, percentage inhibition and  $\text{IC}_{50}$  of compounds under test and significance tests can be calculated. In addition to MCP-1 induced chemotaxis, this alternative form of the assay was also used to measure inhibition of RANTES (2nM) induced chemotaxis.

**d) Binding to human peripheral blood mononuclear cells(PBMCs)**

**i) Preparation of human PBMCs**

Fresh human blood (200ml) was obtained from volunteer donors, collected into sodium citrate anticoagulant to give a final concentration of 0.38%. The blood was mixed with Sedimentation Buffer and incubated at 37°C for 20 minutes. The supernatant was collected and centrifuged at 1700rpm for 5 minutes (Sorvall RT6000D). The pellet obtained was resuspended in 20 ml RPMI/BSA (1mg/ml) and 4 x 5mls of cells were carefully layered over 4 x 5mls of Lymphoprepä (Nycomed) in 15ml centrifuge tubes. Tubes were spun at 1700rpm for 30 minutes (Sorvall RT6000D) and the resultant layer of cells was removed and transferred to 50ml Falcon tubes. The cells were washed twice in Lysis Buffer to remove any remaining red blood cells followed by 2 washes in RPMI/BSA. Cells were resuspended in 5mls of Binding Buffer. Cell number was measured on a Coulter counter and additional binding buffer was added to give a final concentration of  $1.25 \times 10^7$  PBMCs /ml.

**ii) Assay**

[ $^{125}\text{I}$ ]MCP-1 was prepared using Bolton and Hunter conjugation (Bolton *et al.*, 1973, *Biochem. J.*, **133**, 529; Amersham International plc]. Equilibrium binding assays were carried out using the method of Ernst *et al.*, 1994, *J. Immunol.*, **152**, 3541. Briefly, 50 $\mu\text{l}$  of  $^{125}\text{I}$ -labeled MCP-1 (final concentration 100pM) was added to 40 $\mu\text{l}$  ( $5 \times 10^5$  cells) of cell suspension in a 96 well plate. Compounds, diluted in Whole Cell Binding Buffer from a stock solution of 10mM in DMSO were added in a final volume of 5 $\mu\text{l}$  to maintain a constant DMSO concentration in the assay of 5%. Total binding was determined in the absence of compound. Non-specific